Measurement of Aquaporin Z Compatibility with Biomimetic Membranes Using A Developed Membrane Protein Compatibility Evaluation Method

Tingwei Ren[a], Mustafa Erbakan[a,b], Mariusz Grzelakowski[c], Peter Butler[d], Manish Kumar[a]

a. Department of Chemical Engineering, The Pennsylvania State University, University Park, PA 16802; b. Current address: Department of Biosystem Engineering, Bozok University, Yozgat 66000, Turkey; c. Applied Biomimetic, Cincinnati, OH 45237, USA; d. Department of Biomedical Engineering, The Pennsylvania State University, University Park, PA 16802

Cell membrane matrices are mainly composed of lipids and membrane proteins. Membrane proteins, present in the membrane, performing a vast majority of functions including signal reception[1], material transport and biochemical reactions[2,3], that occur at membrane-liquid interfaces. High compatibility between cell membranes and membrane proteins are necessary for living organisms to finish complicated tasks and thrive in the nature[4]. Inspired by living cell systems, researchers are using lipids and lipid-like membrane matrices as models for functional biomimetic membranes development and the number of these membrane matrices are increasing. However, only compatible biomimetic membranes specific to different membrane proteins will provide desired membrane performance[5]. Highly compatible biomimetic membranes have high membrane protein reconstitution density without compromising membrane protein single molecule properties. However, there are few accurate methods for compatibility evaluation[4].

Here, we propose and test a method to evaluate membrane protein compatibility with biomimetic membranes (shown as **Figure**). We use aquaporin (AQP) reconstituted biomimetic membranes (ABMs) as examples to verify the accuracy of this evaluation method. ABM is a widely accepted model system for novel biomimetic membrane research. Previous researchers found that ABMs have higher water transport rate and better salt rejection performance than traditional reverse osmosis membranes[6,7]. However, AQPs compatibility with different biomimetic membranes hasn't been studied systematically. We combined Stopped Flow Light Scattering technology and Fluorescence Correlation Spectroscopy technology for compatibility evaluation[8]. AQPs reconstitution density and single channel water permeability were accurately measured at the single molecule level. By reconstituting AQPs into different ABMs (both block copolymers and lipids), we show that AQP single channel water permeability stays similar in different membranes. However, reconstitution density varies in different materials, and AQPs have higher reconstitution density in block copolymers. We have also implemented this evaluation method on other membrane proteins and artificial water channels[9] and these results will be summarized as well.




Figure: Illustration of developed method for measuring membrane protein compatibility.

References:

1. Broselid, S. et al. G protein-coupled Receptor 30 (GPR30) Forms a Plasma Membrane Complex with Membrane-associated Guanylate Kinases (MAGUKs) and Protein Kinase A-anchoring Protein 5 (AKAP5) That Constitutively Inhibits cAMP Production. Journal of Biological Chemistry 289, 22117-22127,(2014).
2. Parker, M. D., Myers, E. J. & Schelling, J. R. Na+-H+ exchanger-1 (NHE1) regulation in kidney proximal tubule. Cell. Mol. Life Sci. 72, 2061-2074,(2015)
3. Krall, J. F. & Korenman, S. G. Soluble Precursor of Membrane-Associated Protein-Kinase in Uterine Smooth-Muscle Cells. Arch. Biochem. Biophys. 199, 479-487,(1980).

4. Shen, Y.-x., Saboe, P. O., Sines, I. T., Erbakan, M. & Kumar, M. Biomimetic membranes: A review. Journal of Membrane Science 454, 359-381,(2014)
5. Nichols, S. P., Koh, A., Storm, W. L., Shin, J. H. & Schoenfisch, M. H. Biocompatible Materials for Continuous Glucose Monitoring Devices. Chem. Rev. 113, 2528-2549,(2013)
6. Kumar, M., Grzelakowski, M., Zilles, J., Clark, M. & Meier, W. Highly permeable polymeric membranes based on the incorporation of the functional water channel protein Aquaporin Z. Proceedings of the National Academy of Sciences of the United States of America 104, 20719-20724,(2007)
7. Kowal, J., Zhang, X., Dinu, I. A., Palivan, C. G. & Meier, W. Planar Biomimetic Membranes Based on Amphiphilic Block Copolymers. ACS Macro Letters 3, 59-63,(2014)
8. Erbakan, M., Shen, Y.-x., Butler, P. J., Kumar, M. & Curtis, W. R. Molecular cloning, overexpression and characterization of a novel water channel protein from Rhodobacter sphaeroides. PLoS ONE 9, e86830,(2014)
9. Shen, Y. X. et al. Highly permeable artificial water channels that can self-assemble into two-dimensional arrays. Proceedings of the National Academy of Sciences of the United States of America 112, 9810-9815, (2015).